

Research Progress of Cyclophilin A in Pulmonary Infectious Diseases

Juan Chen, Wang Deng*

Department of Respiratory and Critical Care Medicine, Second Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China

**Corresponding author:* Wang Deng, dengwang@hospital.cqmu.edu.cn

Copyright: © 2024 Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), permitting distribution and reproduction in any medium, provided the original work is cited.

Abstract:

Cyclophilin A (CyPA) is an immunophilin with peptidyl-prolyl cis-trans isomerase (PPIase) activity, widely expressed in various tissues and organs, including the lungs. Under inflammatory responses or external cytokine stimulation, CyPA is secreted extracellularly and binds to receptors such as CD147, activating signaling pathways like ERK/NF- κ B. This promotes inflammatory cell chemotaxis and cytokine release, playing a role in the pathophysiological processes of various inflammatory diseases. The expression level of CyPA is positively correlated with the severity of inflammation in pulmonary diseases such as chronic airway inflammation, acute respiratory distress syndrome (ARDS), and COVID-19 pneumonia. Targeting CyPA has been shown to reduce inflammation and improve prognosis. This article reviews the research progress of CyPA in common pulmonary infectious diseases, providing insights into its mechanism of action in such conditions.

Keywords:

Cyclophilin A
Chronic airway inflammation
Acute respiratory distress syndrome
Viral pneumonia

Online publication:

1. Introduction

In 1984, Handschumacher *et al.*^[1] first reported an 18 kD protein derived from bovine thymocytes, which was identified as the intracellular receptor for the immunosuppressant cyclosporin A and formally named Cyclophilin A (CyPA). Subsequently, Fischer *et al.*^[2] discovered that CyPA and peptidylprolyl isomerase A (PPIA) are the same molecule,

possessing peptidyl-prolyl cis-trans isomerase activity. CyPA is expressed in various cell types, such as vascular smooth muscle cells, endothelial cells, epithelial cells, and macrophages, and plays a role in multiple system diseases, including those of the cardiovascular, digestive, urinary, nervous, and rheumatic immune systems^[3-6].

Beyond its intracellular functions, such as protein

folding and transport, CyPA has garnered attention in recent years as a novel inflammatory biomarker [7]. In a mouse model of systemic inflammation induced by a high-fat diet, elevated levels of CyPA and CD147 were detected in multiple organs and tissues [8]. Additionally, numerous studies have confirmed CyPA expression in inflammatory diseases such as vascular inflammation, rheumatoid arthritis, sepsis, and periodontitis [7,9,10]. However, its role in pulmonary infectious diseases remains underexplored.

The lungs, as a common site of infection, are a leading cause of mortality worldwide [11]. Pulmonary infections often result in severe respiratory failure and cardiovascular complications, characterized by high incidence, hospitalization rates, intensive care unit admissions, and mortality. For instance, hospitalized patients with community-acquired pneumonia (CAP) have in-hospital and one-year mortality rates of 6.5% and 30.6%, respectively, while for CAP patients admitted to the ICU, these rates increase to 17% and 47%, respectively [12]. Pulmonary infections significantly impact individual quality of life and impose a substantial economic burden on nations.

Accurately assessing the severity and prognosis of pulmonary infections is therefore critical, and targeted therapies may help address issues such as antibiotic overuse and rising resistance. This review discusses the roles and mechanisms of CyPA in pulmonary infectious diseases, aiming to provide a reference for its potential application in these conditions.

2. Structure and pro-inflammatory mechanisms of CyPA

Cyclophilin A, a member of the immunophilin protein family, is composed of 165 amino acids and contains 8 β -strands and 2 α -helices. It exhibits PPIA activity and is widely expressed in both prokaryotic and eukaryotic organisms [13]. CyPA is involved in various processes such as protein folding, cell proliferation and apoptosis, and cholesterol metabolism [5].

In addition to its catalytic role via PPIA activity, CyPA is secreted extracellularly under inflammatory conditions or external cytokine stimulation, functioning as extracellular CyPA (eCyPA) to mediate inflammatory

responses. It exhibits strong chemotactic effects on lymphocytes, monocytes, and neutrophils [14], primarily through specific binding to CD147 [15,16]. However, the specific interaction groups between CyPA and CD147 remain controversial [5].

Apart from CD147, studies have shown that CyPA enhances the signaling specificity of myeloid cell trigger receptor-2 (TREM2), upregulating the production of pro-inflammatory cytokines, suggesting that TREM2 may be a novel receptor for CyPA [17]. CyPA also plays multiple roles in inflammation by participating in bacterial pathogenesis, such as regulating the formation of host actin cytoskeleton or membrane translocation of bacterial toxins [18].

Moreover, CyPA regulates the transcriptional activity of NF- κ B p65, thereby increasing the production of pro-inflammatory cytokines [19]. For example, in atypical pathogens, the adhesion molecule MgPa secreted by *Mycoplasma genitalium* induces the secretion of eCyPA and its binding to CD147 on urinary epithelial cells, activating the extracellular signal-regulated kinase (ERK) phosphorylation/NF- κ B pathway and mediating the adhesion and invasion of *Mycoplasma genitalium* into human urethral epithelial cells [20,21]. Similar mechanisms have been reported in *Mycoplasma pneumoniae* infections [22].

Additionally, CyPA regulates the degradation of the IL-6 membrane receptor gp130, positively modulating the IL-6 trans-signaling pathway and increasing downstream IL-6 and CyPA expression [23]. It promotes inflammation activation by increasing IL-1 β production and facilitates inflammation resolution by enhancing pro-IL-1 β degradation and IL-1-induced epithelial-mesenchymal transition [24].

In summary, CyPA plays a crucial role in inflammatory responses and is implicated in the pathogenesis of common pulmonary infections.

3. CyPA expression and mechanisms in different pulmonary infectious diseases

3.1. Progress in research on CyPA in chronic airway inflammatory diseases

3.1.1. Asthma

Asthma is one of the most common chronic airway inflammatory diseases, affecting approximately 272 million people worldwide as of 2017 [11]. It is

characterized by acute airway spasms, excessive mucus secretion, and pulmonary inflammation caused by various genetic and environmental factors [25]. Stemmy *et al.* [26] detected persistently elevated expression of eCyPA in the bronchoalveolar lavage fluid of a mouse model of chronic allergic asthma. In a rat model of asthma, treatment with recombinant CyPA protein (rCyPA) significantly reduced airway resistance, with no significant difference compared to traditional asthma medications such as terbutaline and hydrocortisone. *In vitro* experiments showed that rCyPA inhibited the secretion of both Th1 and Th2 cytokines [27].

The role of CyPA in asthma pathogenesis is complex. On one hand, CyPA inhibitors are thought to reduce the expression of inflammatory cells in asthma patients by targeting leukocyte recruitment [26]. On the other hand, since asthma is a Th2 immune response, research indicates that CyPA facilitates asthma development by catalyzing interleukin-2-inducible T-cell kinase (Itk) dimerization, reducing Itk protein kinase activity, and inhibiting T-cell activation [28].

Additionally, oxidative stress is a key mechanism in asthma pathogenesis, and antioxidant therapy has been shown to improve airway remodeling and hyperreactivity [29]. CyPA protects cardiomyocytes from oxidative damage via the AKT/Nox2 pathway [30]. Based on this, researchers suggest that oxidative stress may also be a mechanism by which CyPA contributes to asthma and propose CyPA as a potential therapeutic target for the disease [27].

3.1.2. Chronic obstructive pulmonary disease

Chronic obstructive pulmonary disease (COPD) is another common chronic airway inflammatory disease characterized by irreversible airflow limitation, chronic airway inflammation, and emphysema [31]. In a study of 83 COPD patients with pulmonary infections, CyPA expression in peripheral blood was significantly elevated compared to controls and positively correlated with disease severity, making CyPA a valuable predictor of disease progression in COPD [32].

Another study found significantly increased serum CyPA levels during both the acute exacerbation and recovery phases of COPD. The researchers proposed that CyPA enhances its pro-inflammatory effects by inducing the expression of IL-6 and MMP-9. Serum CyPA levels were negatively correlated with lung function parameters

such as FEV1% and FEV1/FVC during the recovery phase of COPD [33].

These findings suggest that serum CyPA could serve as a potential inflammatory biomarker for COPD, with its levels reflecting disease severity. However, current research on CyPA in COPD is limited to clinical studies, and its molecular mechanisms remain unclear, requiring further investigation.

3.2. Progress in research on CyPA in acute lung injury/acute respiratory distress syndrome

Acute respiratory distress syndrome (ARDS) or acute lung injury is one of the common critical respiratory conditions. It occurs when severe infections, trauma, burns, or other factors cause damage to pulmonary capillary endothelial cells and alveolar epithelial cells, resulting in diffuse pulmonary interstitial and alveolar edema, leading to persistent hypoxemia and respiratory distress. ARDS is considered a primary manifestation of systemic inflammatory response syndrome in the lungs [34].

In a retrospective study, Koh *et al.* [35] analyzed animal and human bronchoalveolar lavage fluid (BALF) samples and proposed that eCyPA is a biomarker for ventilator-induced lung injury (VILI). Compared to healthy volunteers, eCyPA levels in the BALF of ARDS patients were 5–6 times higher. The study also found a positive correlation between eCyPA levels in BALF from VILI mice and markers of lung injury and alveolar epithelial permeability. Blocking CyPA improved survival rates and lung injury, including lung function and oxygenation, in overventilated mice [35]. This suggests the potential value of eCyPA in reflecting alveolar epithelial injury.

In a sepsis mouse model, neutralizing mouse CyPA with an anti-*Clonorchis sinensis* CyPA antibody increased the 72-hour survival rate of septic mice. The protective effect was dose-dependent, with treated mice showing reduced histopathological damage and inflammatory cell infiltration in the lungs and other tissues [36]. This finding supports earlier studies showing that targeting CyPA alleviates lung injury in septic mice [37].

In the pathogenesis of ARDS, experiments have demonstrated that alveolar epithelial cells secrete eCyPA under mechanical stress, such as stretching. eCyPA interacts with CD147 to activate alveolar macrophages,

inducing cytokine secretion and matrix metalloproteinase release within the lungs, ultimately increasing the permeability of the alveolar epithelial barrier^[35].

CD147 plays a significant role in the pathophysiology of acute lung injury, but understanding of the downstream mechanisms activated by CD147 remains limited. Koh *et al.*^[35] also reported that eCyPA levels in serum samples from acute lung injury patients did not differ significantly from controls. Furthermore, early BALF samples are typically unavailable in patients who can breathe independently.

Although CyPA is a promising therapeutic target for mitigating lung injury and improving survival in ARDS patients, clinical application remains challenging. Gaining a better understanding of CyPA's mechanisms of action and developing simpler methods for sample collection will aid in addressing these issues^[38].

3.3. Progress in research on CyPA in viral pneumonia

3.3.1. COVID-19

The coronavirus disease 2019 (COVID-19) is highly infectious and has rapidly spread worldwide since its outbreak in Wuhan in late 2019. Both the number of infections and the geographic spread of the epidemic have far surpassed previous severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS) outbreaks^[39], posing a significant threat to global public health. A meta-analysis of host genes associated with beta-coronavirus infections identified the gene encoding CyPA (PPIA) as the most significant among over 5,000 related genes^[40]. ELISA analyses revealed that CyPA levels in the plasma of severe/critical COVID-19 patients were significantly higher than in mild cases and healthy controls. Additionally, in lung tissues of COVID-19 patients, CyPA was strongly positively correlated with CD68, CCL2, and IL-6^[41]. This suggests that CyPA is linked to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and may serve as a critical pro-inflammatory factor.

A multicenter, randomized, double-blind, controlled international clinical trial showed that the CD147 inhibitor Meplazumab produced beneficial outcomes in severe COVID-19 patients, including reduced mortality, viral load, and cytokine levels^[42]. Animal experiments

confirmed that *CD147* gene knockout significantly inhibited SARS-CoV-2 infection, and Meplazumab effectively suppressed all viral variants except Kappa^[43]. These findings underline the role of CD147 in SARS-CoV-2 infection and suggest its potential as a therapeutic target.

Researchers have proposed the “spike protein-CD147-CyPA signaling axis” as a mechanism for the cytokine storm in COVID-19. Spike protein binding to CD147 activates the JAK-STAT pathway, inducing CyPA expression. CyPA then binds to CD147, triggering the MAPK pathway, which regulates cytokine and chemokine expression, thereby promoting the cytokine storm^[41]. However, some scholars argue that CD147 directly or indirectly influences SARS-CoV-2 entry into host cells via angiotensin-converting enzyme 2 (ACE2), rather than through the CD147-CyPA complex^[44]. Despite these debates, targeting the CyPA/CD147 axis is regarded as a promising therapeutic approach for COVID-19.

3.3.2. Other viral pneumonia

Beyond COVID-19, CyPA plays a role in the pathogenesis of other viral pneumonia. For example, CyPA expression was significantly upregulated in bone marrow-derived macrophages infected with influenza A virus (IAV)^[45]. Bai *et al.*^[46] demonstrated in an IAV-induced mouse model that CyPA promotes integrin $\alpha 5$ expression and actin cytoskeleton rearrangement via the FAK/Akt pathway, enhancing *Streptococcus* group A infection and increasing pulmonary inflammatory infiltration. Further studies revealed that cyclosporin binding to CyPA reduced IAV-induced inflammation by shifting macrophage polarization from the pro-inflammatory M1 phenotype to the anti-inflammatory M2 phenotype through the IFN- γ /STAT1 and IL-4/STAT6 pathways^[47].

Other studies have shown that CyPA interacts with the IAV matrix protein M1, regulating ubiquitin-proteasome-mediated M1 degradation to inhibit viral replication^[48]. In mice infected with influenza B virus (IBV), enzymes promoting CyPA degradation (proteolysis-targeting chimeras, PROTACs) reduced inflammation and lung damage. Combined treatment with oseltamivir and PROTACs was most effective in suppressing inflammatory cytokines^[49]. These findings suggest that targeting CyPA with PROTACs may be a

potential therapeutic strategy for IBV-induced pneumonia.

3.3.3. Balancing pro-inflammatory and antiviral roles

In addition to its pro-inflammatory effects, CyPA also has antiviral properties. For example, human CyPA can block the binding of the SARS-CoV-2 receptor-binding domain (RBD) to the ACE2 receptor, thereby preventing SARS-CoV-2 entry into host cells^[50]. Dittmar *et al.*^[51] demonstrated that cyclosporin analogs exhibit strong anti-SARS-CoV-2 activity in alveolar epithelial cells in a CyPA-dependent manner. In a respiratory syncytial virus (RSV) mouse model, CyPA knockdown resulted in more severe lung inflammation and increased inflammatory cell infiltration compared to controls^[52], suggesting that CyPA inhibits RSV replication.

Therefore, achieving a balance between the pro-inflammatory and antiviral roles of CyPA is crucial for the development of CyPA-targeted therapies for viral pneumonia.

4. Summary

Cyclophilin A is a multifunctional protein that plays a critical role in various pathological conditions. It is secreted from cells through paracrine or autocrine mechanisms, mediating the release of inflammatory cytokines. The expression level of CyPA reflects the severity and prognosis of inflammation. Targeting CyPA for therapy can reduce pulmonary inflammation and improve disease outcomes.

Thus, CyPA can serve as a biomarker for the diagnosis and prognosis of various inflammatory lung diseases and holds promise as a key therapeutic target for alternative treatments in pulmonary infectious diseases. However, current research on CyPA-targeted therapies primarily focuses on its antiviral properties, with clinical applications in anti-inflammatory treatment remaining limited. Additionally, studies on CyPA in bacterial, *Mycobacterium tuberculosis*, and atypical pathogen infections and treatments are relatively scarce, requiring further investigation.

Funding

- (1) Chongqing Municipal Health and Science Joint General Project (Project No. 2023MSXM091)
- (2) Chongqing Science and Technology Bureau General Project (Project No. cstc2020jcyj-msxmX0008)
- (3) Chongqing High-Level Young and Middle-Aged Medical Talent Project

Author contributions

Conceptualization: Wang Deng

Investigation: Juan Chen

Writing – original draft: Juan Chen

Writing – review & editing: Wang Deng

Disclosure statement

The authors declare no conflict of interest.

References

- [1] Handschumacher RE, Harding MW, Rice J, et al., 1984, Cyclophilin: A Specific Cytosolic Binding Protein for Cyclosporin A. *Science*, 226(4674): 544–547. <https://doi.org/10.1126/science.6238408>
- [2] Fischer G, Wittmann-Liebold B, Lang K, et al., 1989, Cyclophilin and Peptidyl-Prolyl Cis-Trans Isomerase are Probably

- Identical Proteins. *Nature*, 337(6206): 476–478. <https://doi.org/10.1038/337476a0>
- [3] Kurosawa R, Satoh K, Nakata T, et al., 2021, Identification of Celastrol as a Novel Therapeutic Agent for Pulmonary Arterial Hypertension and Right Ventricular Failure Through Suppression of Bsg (Basigin)/CyPA (Cyclophilin A). *Arterioscler Thromb Vasc Biol*, 41(3): 1205–1217. <https://doi.org/10.1161/ATVBAHA.120.315731>
- [4] Alvariño R, Alfonso A, Pérez-Fuentes N, et al., 2022, Extracellular Cyclophilins A and C Induce Dysfunction of Pancreatic Microendothelial Cells. *Front Physiol*, 13: 980232. <https://doi.org/10.3389/fphys.2022.980232>
- [5] Hadpech S, Thongboonkerd V, 2022, Current Update on Theranostic Roles of Cyclophilin A in Kidney Diseases. *Theranostics*, 12(9): 4067–4080. <https://doi.org/10.7150/thno.72948>
- [6] Montagne A, Nikolakopoulou AM, Huuskonen MT, et al., 2021, APOE4 Accelerates Advanced-Stage Vascular and Neurodegenerative Disorder in Old Alzheimer’s Mice via Cyclophilin A Independently of Amyloid- β . *Nat Aging*, 1(6): 506–520. <https://doi.org/10.1038/s43587-021-00073-z>. Erratum in *Nat Aging*, 1(7): 624. <https://doi.org/10.1038/s43587-021-00090-y>
- [7] Liao Y, Luo D, Peng K, et al., 2021, Cyclophilin A: A Key Player for Etiological Agent Infection. *Appl Microbiol Biotechnol*, 105(4): 1365–1377. <https://doi.org/10.1007/s00253-021-11115-2>
- [8] Gegunde S, Alfonso A, Cifuentes JM, et al., 2023, Cyclophilins Modify Their Profile Depending on the Organ or Tissue in a Murine Inflammatory Model. *Int Immunopharmacol*, 120: 110351. <https://doi.org/10.1016/j.intimp.2023.110351>
- [9] Yu S, Li J, Yang T, et al., 2022, Expression of CypA/CD147 and MMP-2 in CD68+ Cells of Human Gingival Tissues with Chronic Periodontitis. *Chin J Pathophysiol*, 38(11): 2055–2062.
- [10] Satoh K, Satoh T, Kikuchi N, et al., 2014, Basigin Mediates Pulmonary Hypertension by Promoting Inflammation and Vascular Smooth Muscle Cell Proliferation. *Circ Res*, 115(8): 738–750. <https://doi.org/10.1161/CIRCRESAHA.115.304563>
- [11] GBD 2017 Disease and Injury Incidence and Prevalence Collaborators, 2018, Global, Regional, and National Incidence, Prevalence, and Years Lived with Disability for 354 Diseases and Injuries for 195 Countries and Territories, 1990–2017: A Systematic Analysis for the Global Burden of Disease Study 2017. *Lancet*, 392(10159): 1789–1858. [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7). Erratum in *Lancet*, 393(10190): e44. [https://doi.org/10.1016/S0140-6736\(19\)31047-5](https://doi.org/10.1016/S0140-6736(19)31047-5)
- [12] Cavallazzi R, Ramirez JA, 2022, How and When to Manage Respiratory Infections Out of Hospital. *Eur Respir Rev*, 31(166): 220092. <https://doi.org/10.1183/16000617.0092-2022>
- [13] Favretto F, Flores D, Baker JD, et al., 2020, Catalysis of Proline Isomerization and Molecular Chaperone Activity in a Tug-of-War. *Nat Commun*, 11(1): 6046. <https://doi.org/10.1038/s41467-020-19844-0>
- [14] Dawar FU, Xiong Y, Khattak MNK, et al., 2017, Potential Role of Cyclophilin A in Regulating Cytokine Secretion. *J Leukoc Biol*, 102(4): 989–992. <https://doi.org/10.1189/jlb.3RU0317-090RR>
- [15] Chiu PF, Su SL, Tsai CC, et al., 2018, Cyclophilin A and CD147 Associate with Progression of Diabetic Nephropathy. *Free Radic Res*, 52(11–12): 1456–1463. <https://doi.org/10.1080/10715762.2018.1523545>
- [16] Sakamoto M, Miyagaki T, Kamijo H, et al., 2021, CD147-Cyclophilin A Interactions Promote Proliferation and Survival of Cutaneous T-Cell Lymphoma. *Int J Mol Sci*, 22(15): 7889. <https://doi.org/10.3390/ijms22157889>
- [17] Ji KY, Kim SM, Yee SM, et al., 2021, Cyclophilin A is an Endogenous Ligand for the Triggering Receptor Expressed on Myeloid Cells-2 (TREM2). *FASEB J*, 35(4): e21479. <https://doi.org/10.1096/fj.202002325RR>
- [18] Dhanda AS, Warren KE, Chiu RH, et al., 2018, Cyclophilin A Controls Salmonella Internalization Levels and is Present at E. coli Actin-Rich Pedestals. *Anat Rec (Hoboken)*, 301(12): 2086–2094. <https://doi.org/10.1002/ar.23957>
- [19] Zhai D, Fu Z, Jia J, et al., 2017, Cyclophilin A Aggravates Collagen-Induced Arthritis via Promoting Classically Activated Macrophages. *Inflammation*, 40(5): 1761–1772. <https://doi.org/10.1007/s10753-017-0619-0>
- [20] Deng X, Dai P, Yu M, et al., 2018, Cyclophilin A is the Potential Receptor of the Mycoplasma genitalium Adhesion Protein. *Int J Med Microbiol*, 308(3): 405–412. <https://doi.org/10.1016/j.ijmm.2018.03.001>

- [21] Li L, Luo D, Liao Y, et al., 2020, Mycoplasma genitalium Protein of Adhesion Induces Inflammatory Cytokines via Cyclophilin A-CD147 Activating the ERK-NF- κ B Pathway in Human Urothelial Cells. *Front Immunol*, 11: 2052. <https://doi.org/10.3389/fimmu.2020.02052>
- [22] Reddy SP, Rasmussen WG, Baseman JB, 1996, Correlations between Mycoplasma pneumoniae Sensitivity to Cyclosporin A and Cyclophilin-Mediated Regulation of Mycoplasma Cytoadherence. *Microb Pathog*, 20(3): 155–169. <https://doi.org/10.1006/mpat.1996.0014>
- [23] Luan X, Yang W, Bai X, et al., 2021, Cyclophilin A is a Key Positive and Negative Feedback Regulator Within Interleukin-6 Trans-Signaling Pathway. *FASEB J*, 35(11): e21958. <https://doi.org/10.1096/fj.202101044RRR>
- [24] Yang W, Bai X, Luan X, et al., 2022, Delicate Regulation of IL-1 β -Mediated Inflammation by Cyclophilin A. *Cell Rep*, 38(11): 110513. <https://doi.org/10.1016/j.celrep.2022.110513>. Erratum in *Cell Rep*, 40(12): 111421. <https://doi.org/10.1016/j.celrep.2022.111421>
- [25] Chung KF, Dixey P, Abubakar-Waziri H, et al., 2022, Characteristics, Phenotypes, Mechanisms and Management of Severe Asthma. *Chin Med J (Engl)*, 135(10): 1141–1155. <https://doi.org/10.1097/CM9.0000000000001990>
- [26] Stemmy EJ, Balsley MA, Jurjus RA, et al., 2011, Blocking Cyclophilins in the Chronic Phase of Asthma Reduces the Persistence of Leukocytes and Disease Reactivation. *Am J Respir Cell Mol Biol*, 45(5): 991–998. <https://doi.org/10.1165/rccb.2011-0007OC>
- [27] Chen CT, Shan CX, Ran J, et al., 2021, Cyclophilin A Plays Potential Roles in a Rat Model of Asthma and Suppression of Immune Response. *J Asthma Allergy*, 14: 471–480. <https://doi.org/10.2147/JAA.S308938>
- [28] Schwartzberg PL, Finkelstein LD, Readinger JA, 2005, TEC-Family Kinases: Regulators of T-Helper-Cell Differentiation. *Nat Rev Immunol*, 5(4): 284–295. <https://doi.org/10.1038/nri1591>
- [29] Michaeloudes C, Abubakar-Waziri H, Lakhdar R, et al., 2022, Molecular Mechanisms of Oxidative Stress in Asthma. *Mol Aspects Med*, 85: 101026. <https://doi.org/10.1016/j.mam.2021.101026>
- [30] Cheng F, Yuan W, Cao M, et al., 2019, Cyclophilin A Protects Cardiomyocytes against Hypoxia/Reoxygenation-Induced Apoptosis via the AKT/Nox2 Pathway. *Oxid Med Cell Longev*, 2019: 2717986. <https://doi.org/10.1155/2019/2717986>
- [31] Christenson SA, Smith BM, Bafadhel M, et al., 2022, Chronic Obstructive Pulmonary Disease. *Lancet*, 399(10342): 2227–2242. [https://doi.org/10.1016/S0140-6736\(22\)00470-6](https://doi.org/10.1016/S0140-6736(22)00470-6)
- [32] Liao W, Xu Y, Hu F, et al., 2023, Distribution of Pathogens Isolated from chronic Obstructive Pulmonary Disease Patients with Pulmonary Infection Expressions of CyPA TLR in Peripheral Blood. *Chin J Nosocomiol*, 33(5): 693–697.
- [33] Zhang M, Tang J, Yin J, et al., 2018, The Clinical Implication of Serum Cyclophilin A in Patients with Chronic Obstructive Pulmonary Disease. *Int J Chron Obstruct Pulmon Dis*, 13: 357–363. <https://doi.org/10.2147/COPD.S152898>
- [34] Bos LDJ, Ware LB, 2022, Acute Respiratory Distress Syndrome: Causes, Pathophysiology, and Phenotypes. *Lancet*, 400(10358): 1145–1156. [https://doi.org/10.1016/S0140-6736\(22\)01485-4](https://doi.org/10.1016/S0140-6736(22)01485-4)
- [35] Koh MW, Baldi RF, Soni S, et al., 2021, Secreted Extracellular Cyclophilin A Is a Novel Mediator of Ventilator-induced Lung Injury. *Am J Respir Crit Care Med*, 204(4): 421–430. <https://doi.org/10.1164/rccm.202009-3545OC>
- [36] Jiang J, Yin H, Sun Y, et al., 2018, Clonorchis sinensis Cyclophilin A Immunization Protected Mice from CLP-Induced Sepsis. *Int Immunopharmacol*, 59: 347–353. <https://doi.org/10.1016/j.intimp.2018.03.039>
- [37] Song T, Yang M, Chen J, et al., 2015, Prognosis of Sepsis Induced by Cecal Ligation and Puncture in Mice Improved by Anti-Clonorchis sinensis Cyclophilin A Antibodies. *Parasit Vectors*, 8: 502. <https://doi.org/10.1186/s13071-015-1111-z>
- [38] Witznath M, Kuebler WM, 2021, The CypA-Netics of Ventilator-Induced Lung Injury. *Am J Respir Crit Care Med*, 204(4): 385–387. <https://doi.org/10.1164/rccm.202104-0919ED>
- [39] Hu B, Guo H, Zhou P, et al., 2021, Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol*, 19(3): 141–154. <https://doi.org/10.1038/s41579-020-00459-7>. Erratum in *Nat Rev Microbiol*, 20(5): 315. <https://doi.org/10.1038/s41579-020-00459-7>

022-00711-2

- [40] Parkinson N, Rodgers N, Head Fourman M, et al., 2020, Dynamic Data-Driven Meta-Analysis for Prioritisation of Host Genes Implicated in COVID-19. *Sci Rep*, 10(1): 22303. <https://doi.org/10.1038/s41598-020-79033-3>
- [41] Geng J, Chen L, Yuan Y, et al., 2021, CD147 Antibody Specifically and Effectively Inhibits Infection and Cytokine Storm of SARS-CoV-2 and Its Variants Delta, Alpha, Beta, and Gamma. *Signal Transduct Target Ther*, 6(1): 347. <https://doi.org/10.1038/s41392-021-00760-8>
- [42] Bian H, Chen L, Zheng ZH, et al., 2023, Meplazumab in Hospitalized Adults with Severe COVID-19 (DEFLECT): A Multicenter, Seamless Phase 2/3, Randomized, Third-Party Double-Blind Clinical Trial. *Signal Transduct Target Ther*, 8(1): 46. <https://doi.org/10.1038/s41392-023-01323-9>
- [43] Wang K, Chen W, Zhang Z, et al., 2020, CD147-Spike Protein is a Novel Route for SARS-CoV-2 Infection to Host Cells. *Signal Transduct Target Ther*, 5(1): 283. <https://doi.org/10.1038/s41392-020-00426-x>
- [44] Fenizia C, Galbiati S, Vanetti C, et al., 2021, SARS-CoV-2 Entry: At the Crossroads of CD147 and ACE2. *Cells*, 10(6): 1434. <https://doi.org/10.3390/cells10061434>
- [45] Liu W, Li J, Zheng W, et al., 2017, Cyclophilin A-Regulated Ubiquitination is Critical for RIG-I-Mediated Antiviral Immune Responses. *Elife*, 6: e24425. <https://doi.org/10.7554/eLife.24425>
- [46] Bai X, Yang W, Luan X, et al., 2021, Induction of Cyclophilin A by Influenza A Virus Infection Facilitates Group A Streptococcus Coinfection. *Cell Rep*, 35(7): 109159. <https://doi.org/10.1016/j.celrep.2021.109159>
- [47] Bai X, Yang W, Li H, et al., 2022, Cyclosporine A Regulates Influenza A Virus-induced Macrophages Polarization and Inflammatory Responses by Targeting Cyclophilin A. *Front Immunol*, 13: 861292. <https://doi.org/10.3389/fimmu.2022.861292>
- [48] Mahesutihan M, Zheng W, Cui L, et al., 2018, CypA Regulates AIP4-Mediated M1 Ubiquitination of Influenza A Virus. *Virol Sin*, 33(5): 440–448. <https://doi.org/10.1007/s12250-018-0058-6>
- [49] Li H, Yang W, Li H, et al., 2023, PROTAC Targeting Cyclophilin A Controls Virus-Induced Cytokine Storm. *iScience*, 26(9): 107535. <https://doi.org/10.1016/j.isci.2023.107535>
- [50] Sekhon SS, Shin WR, Kim SY, et al., 2023, Cyclophilin A-Mediated Mitigation of Coronavirus SARS-CoV-2. *Bioeng Transl Med*, 8(2): e10436. <https://doi.org/10.1002/btm2.10436>
- [51] Dittmar M, Lee JS, Whig K, et al., 2021, Drug Repurposing Screens Reveal Cell-Type-Specific Entry Pathways and FDA-Approved Drugs Active Against SARS-Cov-2. *Cell Rep*, 35(1): 108959. <https://doi.org/10.1016/j.celrep.2021.108959>
- [52] Liang W, Zhang Y, Li M, et al., 2021, Cyclophilin A Inhibits Human Respiratory Syncytial Virus (RSV) Replication by Binding to RSV-N through Its PPIase Activity. *J Virol*, 95(15): e0056321. <https://doi.org/10.1128/JVI.00563-21>

Publisher's note

Whioce Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.